

## SELECTIVE SORPTION AND FIXATION OF CATIONS BY CLAY MINERALS: A REVIEW

B. L. SAWHNEY

The Connecticut Agricultural Experiment Station, New Haven, Conn. 06504, U.S.A.

(Received 1 November 1971)

**Abstract**—Investigations concerning selective sorption and fixation of K and similar cations by clay minerals and soil clays and the mechanisms of these reactions are reviewed. In particular, recent observations on selective sorption of these ions in dilute solutions by weathered micas and vermiculite in relation to the interlayer structures are discussed in detail. Also, implications of the resistance to weathering of small mica particles to cation selectivity by soils are described. Despite the increased understanding of sorption and fixation reactions, the following aspects remain unclear.

First, the mechanism of the collapse of alternate layers in vermiculite on K or Cs sorption has not been unequivocally established. Second, factors that impart stability to the central core of mica particles so that K extraction becomes progressively difficult are not known. Third, inability of Ca or Mg ions to expand interlayers of Cs-saturated vermiculite in contrast to K-saturated vermiculite is not completely understood.

### INTRODUCTION

SORPTION of cations from solution by clay minerals and by soils and sediments containing these minerals is usually considered a simple ion exchange process. In non-selective sorption, the amounts of cations sorbed are proportional to their relative concentrations in solution and their sorption can be described by the laws of mass action. Certain cations are, however, sorbed more selectively than others and are held more tightly against replacement by other cations. Potassium,  $\text{NH}_4^+$  and other large monovalent cations are sometimes held so strongly that they are said to be fixed.

Selective sorption of  $\text{K}^+$ ,  $\text{NH}_4^+$ , and  $\text{Cs}^+$  ions by clays has been observed by a number of investigators (Krishnamoorthy and Overstreet, 1950; Wiklander, 1950; Marshall and Garcia, 1959). Numerous early reports on  $\text{K}^+$  fixation have been reviewed by Reitemeier (1951). Although fixation of  $\text{K}^+$  and  $\text{NH}_4^+$  ions by clay minerals had earlier been attributed to their close fit within the hexagonal cavities of basal oxygen planes (Page and Baver, 1940; Barshad, 1948, 1950; Stanford, 1948; Wear and White, 1951), low hydration energy of the ions is now considered to be the major factor in cation selectivity and fixation (Norrish, 1954; Shainberg and Kemper, 1966; Kittrick, 1966). Data showing greater sorption and fixation of  $\text{Cs}^+$  and  $\text{Rb}^+$  ions with smaller hydration energy than of  $\text{K}^+$  ions (Coleman *et al.*, 1963; Sawhney, 1964; Marshall and McDowell, 1965) can be similarly explained.

Cations with low hydration energy, such as

$\text{K}^+$ ,  $\text{NH}_4^+$ ,  $\text{Rb}^+$ , and  $\text{Cs}^+$  produce interlayer dehydration and layer collapse and are therefore fixed in interlayer positions. Conversely, cations with high hydration energy, such as  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{Sr}^{2+}$ , produce expanded interlayers and are not fixed.

The layer charge on the mineral also effects the interlayer collapse and hence, the degree of cation fixation. For instance, in vermiculite,  $\text{K}^+$ -saturation effects interlayer collapse, producing a 10A structure, whereas in montmorillonite with a smaller layer charge than vermiculite,  $\text{K}^+$ -saturation effects only a partial interlayer collapse, producing a 12A structure. Consequently,  $\text{K}^+$  is fixed by vermiculite but  $\text{K}^+$ -saturated montmorillonite must be heated to collapse interlayers to 10A to fix  $\text{K}^+$ . Within the montmorillonite mineral group, the greater the layer charge, the greater is the  $\text{K}^+$  fixation (Weir, 1965). Similar relationships have been observed for  $\text{K}^+$  or  $\text{Cs}^+$  fixation in vermiculite (Sawhney, 1969b). It has been observed also that  $\text{K}^+$  ions are held more tightly in dioctahedral than in trioctahedral minerals (Jackson and Sherman, 1953; Newman and Brown, 1966). Although the exact reasons for this difference are not known, the following two factors may contribute to greater fixation of  $\text{K}^+$  in dioctahedral minerals. First, the vertically oriented OH-dipole of the hydroxyls in trioctahedral layer silicates results in a weaker bonding of  $\text{K}^+$  than the inclined OH-dipole in the dioctahedral layer silicates (Serratos and Bradley, 1958). Second, the smaller size of the octahedral layer in dioctahedral minerals produces shorter

K—O bonds, hence  $K^+$  is held more tightly in dioctahedral minerals (Radoslovich, 1962; Rich, 1968).

#### *Role of frayed edges in selective sorption*

Interest in radioactive fallout and in the disposal of radioactive effluents containing small amounts of  $^{137}Cs^+$  led to investigations that revealed the role of different exchange sites in selective sorption and fixation of  $K^+$  and  $Cs^+$  ions. Results of Jacobs and Tamura (1960) and Tamura and Jacobs (1960) indicated that illite, with a smaller cation exchange capacity than vermiculite, sorbed more Cs than vermiculite from dilute solutions in the range of  $10^{-5}$  M Cs. They attributed this to the selective sorption of Cs in the interlayer space near particle edges. Coleman *et al.* (1963) and Sawhney (1964, 1965) further demonstrated the selective sorption of  $Cs^+$  over  $Ca^{2+}$  and  $Na^+$  ions. Although selective sorption of  $Cs^+$  from dilute solutions was ascribed to the edge sites, the mechanism of selective sorption was not clear.

Evidence for the existence of exchange sites with different selectivities for  $K^+$  ions was obtained by Bolt *et al.* (1963) in experiments on removal of  $K^+$  from a soil illite. These authors recognized three different sites, basal surface, edge-interlayer and interlayer sites. Potassium ions held on basal surfaces and at the edges were exchangeable with other cations. Bolt *et al.* estimated that while the selectivity of basal surfaces of illite for the sorption of  $K^+$  was twice that of  $Ca^{2+}$ , the selectivity of edge interlayer sites was about 500-fold greater. The release of  $K^+$  from interlayer sites was extremely slow, however, and was possibly controlled by a solid- or film-diffusion process as suggested by Mortland and Ellis (1959).

Other investigations have provided additional data illustrating the importance of edge-interlayer sites in selective sorption of  $K^+$  and  $Cs^+$  ions and have provided a possible mechanism for the selectivity of partially weathered micas and vermiculites with collapsed interlayers. Schwertmann (1962a, b) observed preferential sorption of  $K^+$  over  $Ca^{2+}$  ions by a number of soil clays from dilute solutions. Jackson (1963) and Rich (1964) suggested that the selective sorption of  $K^+$  in soils is due to the presence of frayed edges resulting from weathering in micas. Selective sorption of  $K^+$  by soil clays has also been observed by Keay and Wild (1961), Rich and Black (1964) and Dolcater *et al.* (1968). As demonstrated by optical observations in the laboratory, weathering in mica proceeds from the edges inwards (Walker, 1956, 1959; Mortland, 1958; Reed and Scott, 1962; Rausell-Colom *et al.*, 1965; Scott and Smith, 1966; Boyle *et al.*, 1967;

Sawhney and Voigt, 1969). Thus, partially weathered mica (illite) in soils should consist of a collapsed 10A central core and expanded frayed edges. Cations such as  $K^+$ ,  $NH_4^+$ ,  $Rb^+$  and  $Cs^+$  which produce interlayer collapse would be selectively sorbed at the frayed edges to produce more stable collapsed structure similar to that of the central core. Although selective sorption on scrolls and pits on surfaces of large mica particles has been observed using electron microscope (Raman and Jackson, 1964) and electron microprobe (LeRoux *et al.*, 1970), it should be small compared to sorption on frayed edges of weathered micas.

Because the selectivity of frayed edges of weathered micas for  $K^+$  and  $Cs^+$  ions is ascribed to the ease of collapse of frayed edges and because interlayers in vermiculite also collapse on sorption of these ions, relative selectivities of the two minerals have been evaluated. Sawhney (1970) found that illite and micas had larger selectivity than vermiculite at low concentrations of  $K^+$  and  $Cs^+$  ions, presumably due to larger selectivity of frayed-edge sites in illite and mica than interlayer sites in vermiculite. As the concentrations increased, selectivity of illite and micas became less than that of vermiculite. At higher concentrations, as the highly selective sites of frayed edges become saturated with the cation, sorption occurs on non-selective sites in addition to the selective sites; consequently, the overall selectivity of illite and mica decreases. Although selectivity of vermiculite also decreases with increased K or Cs saturation, the decrease is less in vermiculite than in illite and partially weathered micas because of the smaller ratio of the non-selective to selective sites. The decrease in selectivity of illite for  $K^+$  (Bolt *et al.*, 1963; Schouwenburg and Schuffelen, 1963; Tucker, 1967b), of illite and soil clays for  $Cs^+$  (Coleman and LeRoux, 1965; Sawhney, 1970) and of hydrobiotite for  $Rb^+$  (Reichenback, 1968) on increased saturation with the ion is thus explained by sorption on less selective sites in addition to the highly selective frayed edges.

The higher selectivity of frayed edges than the expanded interlayers requires that as frayed edges in mica are increased by removal of the interlayer  $K^+$ , the selectivity for  $K^+$  or  $Cs^+$  should increase. However, when essentially all the interlayer  $K^+$  is removed and the layers expand, the selectivity should decrease. This phenomenon has been demonstrated by LeRoux and Rich (1969), who observed that selectivity of micas for  $Rb^+$  ions first increased on removal of interlayer  $K^+$  and then decreased. Increased sorption of  $Cs^+$  by soil clays of larger CEC (Sawhney and Frink, 1964), presumably due to increased weathering of mica edges, and partially weathered biotite (Sawhney,

1967a) may thus be explained by increased area of the frayed edges.

The ease of collapse of frayed edges resulting in their large selectivity for  $K^+$  or  $Cs^+$  ions may be due to the following forces, as illustrated in Fig. 1 for one interlayer edge, where the forces are resolved into horizontal and vertical components (as shown by arrows). First, the vertical component of the force of attraction due to the  $K^+$  ion in the interlayer position near the frayed edge and the negative layer charge tends to bring the two ends of the frayed edge together as the cation in the forked edge moves out to be replaced by a  $K^+$  ion. Then, the approaching  $K^+$  ion further increases the attraction between the two ends and collapses the edge to produce the stable structure as that of the central core. Consequently,  $K^+$  and other ions with low hydration energy would be selectively sorbed by the frayed edges of the weathered micas. Large hydrated cations, as  $Ca^{2+}$ , on the other hand, tend to keep the edges apart and prevent the formation of a stable structure as that of the core and hence, are neither selectively sorbed nor fixed.

#### *Role of interlayer collapse in vermiculite on $K^+$ or $Cs^+$ sorption*

Sorption of  $K^+$  or  $Cs^+$  ions by montmorillonites and vermiculites produces a collapse in their interlayers. Montmorillonites collapse to give a 12A c-axis spacing on  $K^+$  or  $Cs^+$  saturation while vermiculites give a 10A spacing on saturation with  $K^+$  and 10.8A on saturation with  $Cs^+$ . When progressively larger amounts of these cations are sorbed by these minerals, in the expanded state produced by saturation with  $Ca^{2+}$ , random collapse of interlayers occurs in montmorillonite. In vermiculite,  $K^+$  or  $Cs^+$  sorption causes collapse in alternate layers producing regularly interstratified 10A and 15A layer sequences (Sawhney, 1967b, 1969a). A vermiculite sample from Libby gave a sharp 15A diffraction peak and its higher orders on saturation with  $Ca^{2+}$ . As the interlayer  $Ca^{2+}$  was progressively exchanged by  $K^+$  (or  $Cs^+$ ), a regularly interstratified component with a 25A spacing increased in amount while the 15A component decreased. When approximately half of the interlayer  $Ca^{2+}$  was replaced by  $K^+$  (or  $Cs^+$ ), the entire sample was




Fig. 1. Selective sorption of a  $K^+$  ion in a frayed edge of a weathered mica sheet.

changed to a regularly interstratified mixture. Further sorption of  $K^+$  (or  $Cs^+$ ) produced collapse of the 15A layers within the interstratified mixture until the entire sample was converted to 10A layers. Similar layer collapse in  $Mg^{2+}$ -saturated vermiculite (Rhoades and Coleman, 1967) and in  $K^+$ -depleted micas (Weed and Leonard, 1968) has been observed. Sawhney and Norrish (unpublished) have observed that sorption of successively larger amounts of  $K^+$  produced regular interstratification in vermiculites obtained by removal of almost all interlayer  $K^+$  from biotite, phlogopite and biotite-vermiculite mixtures.

Conversely, the replacement of  $K^+$  in collapsed 10A layers by  $Mg^{2+}$  or  $Ca^{2+}$  produced expansion in alternate layers (Bassett, 1959; Rausell-Colom *et al.*, 1965; Farmer and Wilson, 1970). Bassett suggested that the removal of  $K^+$  from one layer strengthens the K—O bonds in the adjacent layer so that  $K^+$  removal does not occur in the adjacent but occurs in the next layer, effecting expansion of the alternate layers. Farmer and Wilson proposed that the oxidation of the  $Fe^{2+}$  ions in the two silicate layers on either side of a hydrated interlayer should lead to a stronger retention of  $K^+$  in the two adjacent layers and consequently produce regular interstratification.

Interlayer collapse in alternate layers of vermiculite on  $K^+$  sorption should be the reverse of the expansion of alternate layers of mica on  $K^+$  removal. Thus, it has been hypothesized (Sawhney, 1967b; Rhoades and Coleman, 1967) that collapse in one layer of vermiculite on  $K^+$  sorption reduces the charge density of the adjacent layer. Consequently, the bonding energy of  $K^+$  to the silicate sheet of reduced charge is smaller than the hydration energy of the  $Ca^{2+}$  or  $Mg^{2+}$  ions, hence  $K^+$  would not enter this layer but would enter the next layer, producing regularly interstratified layer sequences. Weed and Leonard (1968) ascribed the prevention of  $K^+$  entry to greater hydration of the layer of reduced charge.

Although these hypotheses have been advanced to explain the expansion of alternate layers in mica on  $K^+$  removal and the collapse of alternate layers in vermiculite on  $K^+$  sorption, no comprehensive treatment of these changes has emerged yet and the mechanism of regular interstratification remains unclear.

#### *Comparison of fixation of $K^+$ and $Cs^+$ by clays*

Just as the fixation of  $K^+$  and  $Cs^+$  ions is accompanied by interlayer collapse, the release of these ions should occur by expansion of the interlayers. Thus, when micas and  $K^+$ -saturated vermiculite are exhaustively leached with solutions of cations, such as  $Ca^{2+}$  and  $Mg^{2+}$ , that produce interlayer

expansion,  $K^+$  is replaced by these cations. Conversely,  $K^+$  is not replaced effectively by cations, such as  $NH_4^+$ , that produce interlayer collapse in minerals (Barshad, 1948, 1950). Indeed,  $K^+$  retained against replacement by  $NH_4^+$  has been commonly used as a measure of  $K^+$  fixation. Furthermore, the presence of small amounts of ions such as  $NH_4^+$  in solutions of  $Ca^{2+}$  and  $Mg^{2+}$  salts used for extracting  $K^+$  inhibits the release of  $K^+$  (Scott and Smith, 1966; Wells and Norrish, 1968).

Recent observations, however, revealed that in contrast to  $K^+$  fixed in the interlayers of vermiculite, a portion of the  $K^+$  in illite and certain soils is more readily replaced by  $NH_4^+$  than by  $Ca^{2+}$  or  $Mg^{2+}$  ions (Bolt *et al.*, 1963; Rich and Black, 1964; Tucker, 1967a). As discussed above,  $K^+$  fixed by illite and weathered micas in soil clays occupies edge-interlayer sites that are highly specific for  $K^+$  and other similar cations. Because of the similarity of  $K^+$  and  $NH_4^+$  ions and the small diffusion distance from solution to these sites,  $NH_4^+$  ions are effective in replacing  $K^+$ . Conversely, hydrated ions  $Ca^{2+}$  and  $Mg^{2+}$  are not selectively sorbed by these sites, hence they are not as effective in replacing  $K^+$ .

Fixation of  $Cs^+$  by layer silicates appears to be similar to that of  $K^+$  at the edge-interlayer sites of partially weathered micas and not as  $K^+$  fixation in interlayers of vermiculite or montmorillonite. Several investigations show that  $K^+$  and  $NH_4^+$  ions replace  $Cs^+$  fixed at the edge-interlayer sites as well as in the interlayer positions more readily than  $Ca^{2+}$  and  $Mg^{2+}$  ions (Schulz *et al.*, 1960; Coleman *et al.*, 1963; Sawhney, 1964). Similarly,  $Cs^+$  fixed in soils and soil clays was replaced more readily by  $K^+$  than  $Ca^{2+}$  (Klechkovsky *et al.*, 1959; Schulz *et al.*, 1960; Nishita *et al.*, 1962; Coleman and LeRoux, 1965). Cesium ions with a smaller hydration energy than  $K^+$  ions hold the layers together more strongly than  $K^+$  ions so that hydrated  $Ca^{2+}$  and  $Mg^{2+}$  ions which can expand layers held by  $K^+$  ions are ineffective in producing layer expansion in  $Cs^+$ -saturated minerals. Consequently,  $Cs^+$  ions are not as readily replaced by these ions as are the  $K^+$  ions. Smaller retention of  $Cs^+$  against replacement by  $K^+$  than  $Ca^{2+}$  or  $Mg^{2+}$  ions has been suggested to be related to interlayer distances (Coleman *et al.*, 1963). For example,  $Cs^+$  fixed by vermiculite or heated montmorillonite produces interlayer distances (basal spacing = 10.8 Å) large enough to permit  $K^+$  ions to diffuse into the interlayers and replace the  $Cs^+$  ions but greatly restrict the entry of hydrated  $Ca^{2+}$  or  $Mg^{2+}$  ions. Although these hypotheses appear logical, lack of precise measurements of the bonding energies of different cations and of the forces involved in interlayer

collapse and expansion preclude an unequivocal explanation for the differences in the interlayer fixation of  $K^+$  and  $Cs^+$  ions.

#### *Release of $K^+$ from naturally-occurring micas and implication in cation selectivity and fixation*

Most of the  $K^+$  in naturally-occurring hydrothermal micas is fixed in the interlayer positions and is not exchangeable, while a small portion that is present on basal surfaces is exchangeable with other cations. When micas are weathered artificially by treatment with a neutral salt solution, interlayer  $K^+$  is slowly replaced by the cation of the neutral salt and the interlayers are expanded to form vermiculite. It has been suggested that in addition to the removal of  $K^+$ , the net negative charge on the layers also decreases during weathering (Jackson and Sherman, 1953; Raman and Jackson, 1966; Newman and Brown, 1966). Newman and Brown concluded that the changes in chemical composition of micas during weathering include the release of structural OH ions exposed on replacement of  $K^+$  or the sorption of  $H^+$  ions and oxidation of iron in the octahedral sheet or loss of divalent cations, resulting in the decrease in net negative charge. Brown and Newman (1970) observed that the removal of interlayer  $K^+$  from micas also produces slight structural irregularities.

Studies of  $K^+$  release in relation to particle size of micas (Mortland and Lawton, 1961; Reichenbach and Rich, 1969; Scott, 1968) show that initial release of  $K^+$  from surface sites is somewhat greater in smaller particles, presumably due to the larger surface area and more surface  $K^+$  on smaller particles. However, the total  $K^+$  released decreases with decreasing particle size. Reichenbach and Rich showed that whereas extraction with  $BaCl_2$  solution over a 5 day period replaced only about 60 me  $K^+$  per 100g from less than 0.08  $\mu$  particles, the same treatment replaced about 200 me per 100g from 5 to 2  $\mu$  particles. Scott also concluded that smaller particles resist the release of  $K^+$ ; over 95% of  $K^+$  was extracted from 60 to 50  $\mu$  particles in 3 yr, using the extraction procedure of Scott and Smith (1966), but less than 70% of  $K^+$  was extracted from 0.7 to 0.2  $\mu$  particles. Similarly, in our experiments (Sawhney and Norrish, unpublished), we found that after a 4-week treatment with  $BaCl_2$  (Rich, 1968; slightly modified), 50–5  $\mu$  particles of biotite and phlogopite retained about 2% and 1%  $K_2O$  respectively. An illite sample (about 1  $\mu$  particle size) retained as much as 7.5%  $K_2O$  following the same treatment.

It appears that removal of interlayer K from micas proceeds from the edges inwards, leaving a stable central core as in Fig. 1. The amount of  $K^+$  retained per unit weight is larger in smaller parti-

cles, as shown in Fig. 2 (data taken from Scott, 1968; Fig. 7). However, the number of  $K^+$  ions retained per particle, and hence the size of the interlayer core increases with increasing particle size, as illustrated in Fig. 2. The diameter of the core was calculated from data of Scott (1968, Fig. 7), and was based on the assumptions that the particles are spherical, they initially contain 250 me K/100g, and each  $K^+$  ion occupies  $450\text{\AA}^3$  of interlayer space of dimensions  $a = 5\text{\AA}$ ,  $b = 9\text{\AA}$  and  $c = 10\text{\AA}$ . Over the range of particle sizes examined, the diameter of the central core does not appear to reach an upper limit. Extrapolating the data downwards to the point where the diameter of the particle and of the core are equal, it appears that the central core becomes extremely stable at a particle diameter of about  $100\text{\AA}$ . The changes in configuration of electrical field which impart stability to the central core are not known, although the resistance of small particles to  $K^+$  release may be related to better structural order in smaller particles. Smaller particles of kaolinite have been shown to be more highly ordered (Ormsby, *et al.*, 1962; Wiewiora and Brindley, 1969), though a similar relationship for illite has not been reported.

The resistance of small particles to release  $K^+$  has important implications in cation selectivity and fixation by soil clays. Calculations from the data of Reichenbach and Rich (1969) and Scott (1968) on  $K^+$  release by different particle size of micas reveal that the area of collapsed central core relative to the frayed edges is larger in the smaller particles. This should impart greater ease of collapse to the frayed edges in smaller particles. Con-

sequently, smaller particles would show greater selectivity for  $K^+$  and similar ions, although the total capacity for fixation may be greater in larger weathered particles (Hill and Sawhney, 1969) due to greater  $K^+$ -depletion in them during weathering. The large selectivity of soils for  $K^+$  in dilute solutions is, thus, attributed to the presence of fine partially weathered mica particles, illite, in soil clays.

Although the investigations reviewed here have described the selective sorption and fixation of  $K^+$  and  $Cs^+$  ions from dilute solutions, the following aspects of the reactions of these ions remain unanswered. First, the mechanism of the collapse of alternate layers in vermiculite on  $K^+$  or  $Cs^+$  sorption has not been unequivocally established. Second, factors that impart stability to the central core of mica particles so that  $K^+$  extraction becomes progressively difficult are not known. And third, the inability of  $Ca^{2+}$  or  $Mg^{2+}$  ions to expand interlayers of  $Cs^+$ -saturated vermiculite in contrast to  $K^+$ -saturated vermiculite is not completely understood.

*Acknowledgments*—This study was supported in part by the U.S. Atomic Energy Commission under contract AT(30-1)-2955.

## REFERENCES

- Barshad, I. (1948) Vermiculite and its relation to biotite as revealed by base exchange reactions, X-ray analyses, differential thermal curves, and water content: *Am. Mineralogist* **33**, 655-678.
- Barshad, I. (1950) The effect of the interlayer cations on the expansion of the mica type of crystal lattice: *Am. Mineralogist* **35**, 225-238.
- Bassett, W. A. (1959) The origin of vermiculite deposit at Libby, Montana: *Am. Mineralogist* **44**, 282-299.
- Bolt, G. H., Sumner, M. E. and Kamphorst, A. (1963) A study of the equilibria between three categories of potassium in an illitic soil: *Soil Sci. Soc. Am. Proc.* **27**, 294-299.
- Boyle, J. R., Voigt, G. K. and Sawhney, B. L. (1967) Biotite flakes: Alteration by chemical and biological treatment: *Science* **155**, 193-195.
- Brown, G. and Newman, A. C. D. (1970) Cation exchange properties of micas—III. Release of potassium sorbed by potassium-depleted micas: *Clay Minerals* **8**, 273-278.
- Coleman, N. T., Craig, Doris, and Lewis, R. J. (1963) Ion-exchange reactions of cesium: *Soil Sci. Soc. Am. Proc.* **27**, 287-289.
- Coleman, N. T. and LeRoux, F. H. (1965) Ion-exchange displacement of cesium from soil vermiculite: *Soil Sci.* **99**, 243-250.
- Dolcater, D. L., Lotse, E. G., Syers, J. K. and Jackson, M. L. (1968) Cation exchange selectivity of some clay-sized minerals and soil materials: *Soil Sci. Soc. Am. Proc.* **32**, 795-798.
- Farmer, V. C. and Wilson, M. J. (1970) Experimental




Fig. 2. Relationship among particle size, diameter of collapsed central core and K retained by muscovite (calculated from data of Scott, 1968).

- conversion of biotite to hydrobiotite: *Nature* **226**, 841-842.
- Hill, D. E. and Sawhney, B. L. (1969) Electron microprobe analysis of thin sections of soils to observe loci of cation exchange: *Soil Sci. Soc. Am. Proc.* **33**, 531-534.
- Jacobs, D. G. and Tamura, T. (1960) The mechanism of ion fixation using radio-isotope techniques: *Trans. Int. Congr. Soil Sci.* 7th Madison **2**, 206-214.
- Jackson, M. L. and Sherman, G. D. (1953) Chemical weathering of minerals in soils: *Advan. Agron.* **5**, 219-318.
- Jackson, M. L. (1963) Interlayering of expansible layer silicates in soils by chemical weathering: *Clays and Clay Minerals* **11**, 29-46.
- Keay, J. and Wild, A. (1961) The kinetics of cation exchange in vermiculite: *Soil Sci.* **92**, 54-60.
- Kittrick, J. A. (1966) Forces involved in ion fixation by vermiculite: *Soil Sci. Soc. Am. Proc.* **30**, 801-803.
- Krishnamoorthy, C. and Overstreet, R. (1950) An experimental evaluation of ion exchange relationships: *Soil Sci.* **69**, 41-53.
- Klechkovsky, V. M., Sokolova, L. N. and Tselishcheva, G. N. (1959) The sorption of microquantities of strontium and cesium in soils: In *Progress in Nuclear Energy* (Edited by W. G. Marley and K. Z. Morgan), Vol. 12, pp. 486-499, *Health Physics*, Pergamon Press, New York.
- LeRoux, J. and Rich, C. I. (1969) Ion selectivity of micas as influenced by degree of potassium depletion: *Soil Sci. Soc. Am. Proc.* **33**, 684-690.
- LeRoux, J., Rich, C. I. and Ribbe, P. H. (1970) Ion selectivity by weathered micas as determined by electron microprobe analysis: *Clays and Clay Minerals* **18**, 333-338.
- Marshall, C. E. and Garcia, G. (1959) Exchange equilibria in a carboxylic resin and in Attapulgite clay: *J. Phys. Chem.* **63**, 1663-1666.
- Marshall, C. E. and McDowell, L. L. (1965) The surface reactivity of micas: *Soil Sci.* **99**, 115-131.
- Mortland, M. M. (1958) Kinetics of potassium release from biotite: *Soil Sci. Soc. Am. Proc.* **22**, 503-508.
- Mortland, M. M. and Ellis, B. G. (1959) Release of fixed potassium as a diffusion controlled process: *Soil Sci. Soc. Am. Proc.* **23**, 363-364.
- Mortland, M. M. and Lawton, K. (1961) Relationships between particle size and potassium release from biotite and its analogues: *Soil Sci. Soc. Am. Proc.* **25**, 473-476.
- Newman, A. C. D. and Brown, G. (1966) Chemical changes during the alteration of micas: *Clay Minerals* **6**, 297-310.
- Nishita, H., Taylor, P., Alexander, G. V. and Larson, K. H. (1962) Influence of stable Cs and K on the reactions of <sup>137</sup>Cs and <sup>42</sup>K in soils and clay minerals: *Soil Sci.* **94**, 187-197.
- Norrish, K. (1954) The swelling of montmorillonite: *Disc. Faraday Soc.* **18**, 120-134.
- Ormsby, W. C., Scharf, J. M. and Woodside, K. H. (1962) Exchange behaviour of kaolins of varying degrees of crystallinity: *J. Am. Ceram. Soc.* **45**, 361-366.
- Page, J. B. and Bayer, L. D. (1940) Ionic size in relation to fixation of cations by colloidal clay: *Soil Sci. Soc. Am. Proc.* **4**, 150-155.
- Radoslovich, E. W. (1962) The cell dimensions and symmetry of layer lattice silicates—II. Regression relations: *Am. Mineralogist* **47**, 617-636.
- Raman, K. V. and Jackson, M. L. (1964) Vermiculite surface morphology: *Clays and Clay Minerals* **12**, 423-429.
- Raman, K. V. and Jackson, M. L. (1966) Layer charge relations in clay minerals of micaceous soils and sediments: *Clays and Clay Minerals* **14**, 53-68.
- Rausell-Colom, J. A., Sweatman, T. R., Wells, C. B. and Norrish, K. (1965) Studies in the artificial weathering of mica: In *Experimental Pedology* (Edited by E. G. Hallsworth and D. V. Crawford), pp. 40-72, Butterworths London.
- Reed, M. G. and Scott, A. D. (1962) Kinetics of potassium release from biotite and muscovite in sodium tetraphenylboron solutions: *Soil Sci. Soc. Am. Proc.* **26**, 437-440.
- Rhoades, J. D. and Coleman, N. T. (1967) Interstratification in vermiculite and biotite produced by potassium sorption—I. Evaluation by simple X-ray diffraction pattern inspection: *Soil Sci. Soc. Am. Proc.* **31**, 366-372.
- Reichenbach, H. G. Von (1968) Cation exchange in the interlayers of expansible layer silicates: *Clay Minerals* **7**, 331-341.
- Reichenbach, H. G. Von and Rich, C. I. (1969) Potassium release from muscovite as influenced by particle size: *Clays and Clay Minerals* **17**, 23-29.
- Reitemeier, R. F. (1951) The chemistry of soil potassium: *Advan. Agron.* **3**, 113-164.
- Rich, C. I. (1964) Effect of cation size and pH on potassium exchange in Nason soil: *Soil Sci.* **98**, 100-106.
- Rich, C. I. and Black, W. R. (1964) Potassium exchange as affected by cation size, pH, and mineral structure: *Soil Sci.* **97**, 384-390.
- Rich, C. I. (1968) Mineralogy of soil potassium: In *The Role of Potassium in Agriculture* (Edited by V. J. Kilmer, S. E. Younts and N. C. Brady), pp. 79-108, American Society of Agronomy, Inc., Madison, Wisc.
- Sawhney, B. L. (1964) Sorption and fixation of microquantities of Cs by clay minerals: effect of saturating cations: *Soil Sci. Soc. Am. Proc.* **28**, 183-186.
- Sawhney, B. L. and Frink, C. R. (1964) Sorption of cesium from dilute solutions by soil clays: *Trans. 8th Int. Cong. Soil Sci. Bucharest* **3**, 423-431.
- Sawhney, B. L. (1965) Sorption of cesium from dilute solutions: *Soil Sci. Soc. Am. Proc.* **29**, 25-28.
- Sawhney, B. L. (1967a) Cesium sorption in relation to lattice spacing and cation exchange capacity of biotite: *Soil Sci. Soc. Am. Proc.* **31**, 181-183.
- Sawhney, B. L. (1967b) Interstratification in vermiculite: *Clays and Clay Minerals* **15**, 75-84.
- Sawhney, B. L. (1969a) Regularity of interstratification as affected by charge density in layer silicates: *Soil Sci. Soc. Am. Proc.* **33**, 42-46.
- Sawhney, B. L. (1969b) Cesium uptake by layer silicates: effect on interlayer collapse and cation exchange capacity: *Proc. Int. Clay Conf.*, Tokyo **1**, 605-611. Israel Universities Press.
- Sawhney, B. L. (1970) Potassium and cesium ion selectivity in relation to clay mineral structure: *Clays and Clay Minerals* **18**, 47-52.
- Sawhney, B. L. and Voigt, G. K. (1969) Chemical and

- biological weathering in vermiculite from Transvaal: *Soil Sci. Soc. Am. Proc.* **33**, 625–629.
- Schulz, R. K., Overstreet, R. and Barshad, I. (1960) On the soil chemistry of  $^{137}\text{Cs}$ : *Soil Sci.* **89**, 16–27.
- Schwertmann, U. (1962a) Eigenschaften und Bildung aufweitbarer (quellbarer) Dreischicht-Tonminerale in Boden aus Sedimenten: *Beitrage zur Mineralogie und Petrographie* **8**, 199–209.
- Schwertmann, U. (1962b) Die selective Kationen sorption der Tonfraktion einiger Boden aus Sedimenten: *Z. Pflanzenernahr. Dung. Bodenk.* **97**, 9–25.
- Shouwenburg, J. C. H. Von and Schuffelen, A. C. (1963) Potassium exchange behaviour of an illite: *Neth. J. Agric. Sci.* **11**, 13–22.
- Scott, A. D. and Smith, S. J. (1966) Susceptibility of interlayer potassium in micas to exchange with sodium: *Clays and Clay Minerals* **14**, 69–81.
- Scott, A. D. (1968) Effect of particle size on interlayer potassium exchange in micas: *Trans. Int. Cong. Soil Sci. Adelaide* **2**, 649–659.
- Serratoso, J. M. and Bradley, W. F. (1958) Determination of the orientation of OH bond axes in layer silicates by infrared adsorption: *J. Phys. Chem.* **62**, 1164–1167.
- Shainberg, I. and Kemper, W. D. (1966) Hydration status of adsorbed cations: *Soil Sci. Soc. Am. Proc.* **30**, 707–713.
- Stanford, G. (1948) Fixation of potassium in soils under moist conditions and on drying in relation to type of clay mineral: *Soil Sci. Soc. Am. Proc.* **12**, 167–171.
- Tamura, T. and Jacobs, D. G. (1960) Structural implications in cesium sorption: *Health Physics* **2**, 391–398.
- Tucker, B. M. (1967a) The solubility of potassium from soil illites—III. Reactivity towards other ions: *Aust. J. Soil Res.* **5**, 173–190.
- Tucker, B. M. (1967b) The solubility of potassium from soil illites—IV. Rate of reaction and exchange constants: *Aust. J. Soil Res.* **5**, 191–201.
- Walker, G. F. (1956) The diffusion of interlayer water in vermiculite: *Nature* **177**, 239–240.
- Walker, G. F. (1959) Diffusion of exchangeable cations in vermiculite: *Nature* **184**, 1392–1394.
- Wear, J. I. and White, J. L. (1951) Potassium fixation in clay minerals as related to crystal structure: *Soil Sci.* **71**, 1–14.
- Weed, S. B. and Leonard, R. A. (1968) Effect of  $\text{K}^+$ -uptake by  $\text{K}^+$ -depleted micas on the basal spacing: *Soil Sci. Soc. Am. Proc.* **32**, 335–340.
- Weir, A. H. (1965) Potassium retention in montmorillonite: *Clay Minerals* **6**, 17–22.
- Wells, C. B. and Norrish, K. (1968) Accelerated rates of release of interlayer potassium from micas: *Trans. Int. Congr. Soil Sci. 9th, Adelaide* **2**, 683–694.
- Wiewiora, A. and Brindley, G. W. (1969) Potassium acetate intercalation in kaolinite and its removal; effect of material characteristics: *Proc. Int. Clay Conf., Tokyo* **1**, 723–733. Israel Universities Press.
- Wiklander, L. (1950) Fixation of potassium by clays saturated with different cations: *Soil Sci.* **69**, 261–268.

**Résumé**—On passe en revue les recherches concernant la sorption sélective et la fixation de K et de cations similaires par les minéraux argileux et les argiles des sols ainsi que les recherches concernant les mécanismes de ces réactions. En particulier, des observations récentes sur la sorption sélective de ces ions en solutions diluées par les micas altérés et la vermiculite, en liaison avec les structures interfeuillettes, sont discutées en détail. De même, on décrit les implications de la résistance à l'altération de petites particules de mica dans la sélectivité des sols pour le cation. En dépit d'une amélioration dans la compréhension des réactions de sorption et de fixation, les points suivants sont toujours mal éclaircis.

En premier lieu, le mécanisme de la fermeture de couches alternées dans la vermiculite lors de la sorption de K ou Cs n'a pas été établi sans équivoque. En second lieu, les facteurs qui confèrent la stabilité au noyau central des particules de mica, si bien que l'extraction de K devient de plus en plus difficile, ne sont pas connus. En troisième lieu, l'inaptitude des ions Ca ou Mg à ouvrir les espaces interfeuillettes de vermiculite saturée par le Cs, ce qui contraste avec le comportement de la vermiculite saturée par le K, n'est pas complètement comprise.

**Kurzreferat**—Es werden Untersuchungen über die selektive Sorption und Fixierung von K und ähnlichen Kationen durch Tonminerale und Bodentone sowie die Mechanismen dieser Reaktionen überprüft. Insbesondere werden neuere Beobachtungen über selektive Sorption dieser Ionen in verdünnten Lösungen durch verwitterte Glimmer und Vermiculite in bezug auf die Zwischenschichtgefüge im Einzelnen erörtert. Ferner wird die Bedeutung des Widerstandes gegen Verwitterung kleiner Glimmerteilchen für die Kationenselektivität der Böden beschrieben. Ungeachtet des besseren Verständnisses der Sorptions- und Fixierungsreaktionen, sind die folgenden Aspekte weiter unklar.

Erstens ist der Mechanismus des Zusammenbruchs alternierender Schichten im Vermiculit bei K oder Cs Sorption nicht eindeutig festgelegt worden. Zweitens sind die Faktoren, die dem inneren Kern von Glimmerteilchen Stabilität verleihen, so dass die K-Extraktion fortschreitend schwieriger wird nicht bekannt. Drittens ist die Unfähigkeit von Ca oder Mg Ionen die Zwischenschichten von Cs-gesättigtem Vermiculit im Gegensatz zum K-gesättigten Vermiculit auszuweiten nicht ganz verständlich.

**Резюме** — Дан обзор исследований, посвященных селективному поглощению и фиксации K и аналогичных катионов глинистыми минералами и почвенными глинами, а также механизму подобных реакций. В частности, детально рассматриваются недавно проведенные исследо-

вания по селективному поглощению этих катионов в разбавленных растворах выветрелыми слюдами и вермикулитом с разной структурой межслоевого пространства. Стойкость к выветриванию небольших частиц слюды объясняется селективностью почв по отношению к катионам. Однако отмечается, что несмотря на все углубляющееся понимание механизма поглощения и фиксации остаются неясными следующие аспекты.

Во-первых, однозначно не установлен механизм сжатия чередующихся слоев в вермикулите в результате поглощения К или Са. Во-вторых, до сих пор не выяснены факторы, определяющие устойчивость центральных частей глинистых частиц, благодаря которым удаление К становится все более и более затруднительным. В-третьих, полностью не понята неспособность ионов Са или Mg вызывать разбухание межслоевых промежутков Cs-насыщенных вермикулитов в противоположность их действию на К-насыщенные вермикулиты.